

A change of wavelength in light scattering

Further observations by Mr Krishnan and myself on the new kind of light-scattering discovered by us have been made and have led to some very surprising and interesting results.

In order to convince ourselves that the secondary radiation observed by us was a true scattering and not a fluorescence, we proceeded to examine the effect in greater detail. The principal difficulty in observing the effect with gases and vapours was its excessive feebleness. In the case of substances of sufficient light-scattering power, this difficulty was overcome by using an enclosed bulb and heating it up so as to secure an adequate density of vapour. Using a blue-violet filter in the track of the incident light, and a complementary green-yellow filter in front of the observer's eye, the modified scattered radiation was observed with a number of organic vapours, and it was even possible to determine its state of polarisation. It was found that in certain cases, for e.g. pentane, it was strongly polarised, while in others, as for example naphthalene, it was only feebly so, the behaviour being parallel to that observed in the liquid state. Liquid carbon dioxide in a steel observation vessel was studied, and exhibited the modified scattering to a notable extent. When a cloud was formed within the vessel by expansion, the modified scattering brightened up at the same time as the ordinary or classical scattering. The conclusion is thus reached that the radiations of altered wavelength from neighbouring molecules are coherent with each other.

A greater surprise was provided by the spectroscopic observations. Using sunlight with a blue filter as the illuminant, the modified scattered radiation was readily detected by the appearance in the spectrum of the scattered light of radiations absent from the incident light. With a suitably chosen filter in the incident light, the classical and modified scatterings appeared as separate regions in the spectrum separated by a dark region. This encouraged us to use a mercury arc as the source of light, all radiations of longer wavelength than 4358 Å being cut out by a filter. The scattered radiations when examined with a spectroscope showed some sharp bright lines additional to those present in the incident light, their wavelength being longer than 4358 Å; at least two such lines were prominent and appeared to be accompanied by some fainter lines, and in addition a continuous spectrum. The relation of frequencies between the new lines and those present in the incident light is being investigated by photographing and measuring the spectra. The preliminary visual observations appear to indicate that the position of the principal modified lines is the same for all substances,

though their intensity and that of the continuous spectrum does vary with their chemical nature.

C V RAMAN

210 Bowbazaar Street, Calcutta
8 March