

2.4. Isomorphous replacement and anomalous scattering

By M. VIJAYAN AND S. RAMASESHAN

2.4.1. Introduction	264	2.4.5. Anomalous scattering of neutrons and synchrotron radiation. The multiwavelength method	275
2.4.2. Isomorphous replacement method	264		
2.4.3. Anomalous-scattering method	266		
2.4.4. Isomorphous replacement and anomalous scattering in protein crystallography	269		

2.4.1. Introduction

Isomorphous replacement is among the earliest methods to be employed for crystal structure determination (Cork, 1927). The power of this method was amply demonstrated in the classical X-ray work of J. M. Robertson on phthalocyanine in the 1930's using centric data (Robertson, 1936; Robertson & Woodward, 1937). The structure determination of strychnine sulfate pentahydrate by Bijvoet and others provides an early example of the application of this method to acentric reflections (Bokhoven, Schoone & Bijvoet, 1951). The usefulness of isomorphous replacement in the analysis of complex protein structures was demonstrated by Perutz and colleagues (Green, Ingram & Perutz, 1954). This was closely followed by developments in the methodology for the application of isomorphous replacement to protein work (Harker, 1956; Blow & Crick, 1959) and rapidly led to the first ever structure solution of two related protein crystals, namely, those of **myoglobin and haemoglobin (Kendrew, Dickerson, Strandberg, Hart, Phillips & Shore, 1960; Cullis, Muirhead, Perutz, Rossmann & North, 1961b)**. Since then isomorphous replacement has been the method of choice in macromolecular crystallography and most of the subsequent developments in and applications of this method have been concerned with biological macromolecules, mainly proteins (Blundell & Johnson, 1976; McPherson, 1982).

The application of anomalous-scattering effects has often developed in parallel with that of isomorphous replacement. Indeed, the two methods are complementary to a substantial extent and they are often treated together, as in this article. Although the most important effect of anomalous scattering, namely, the violation of Friedel's law, was experimentally observed as early as 1930 (Coster, Knol & Prins, 1930), two decades elapsed before this effect was made use of for the first time by Bijvoet and his associates for the determination of the absolute configuration of asymmetric molecules as well as for phase evaluation (Bijvoet, 1949, 1954; Bijvoet, Peerdeman & van Bommel, 1951). Since then there has been a phenomenal spurt in the application of anomalous-scattering effects (Srinivasan, 1972; Ramaseshan & Abrahams, 1975; Vijayan, 1987). A quantitative formulation for the determination of phase angles using intensity differences between Friedel equivalents was derived by Ramachandran & Raman (1956), while Okaya & Pepinsky (1956) successfully developed a Patterson approach involving anomalous effects. The anomalous-scattering method of phase determination has since been used in the structure analysis of several structures, including those of a complex derivative of vitamin B₁₂ (Dale, Hodgkin & Venkatesan, 1963) and a small protein (Hendrickson & Teeter, 1981). In the meantime, the effect of changes in the real component of the dispersion correction as a function of the wavelength of the radiation used, first demonstrated by Mark & Szillard (1925), also received considerable attention. This effect, which is formally equivalent to that of isomorphous replacement, was demonstrated to be useful in structure determination (Ramaseshan, Venkatesan & Mani, 1957;

Ramaseshan, 1963). Protein crystallographers have been quick to exploit anomalous-scattering effects (Rossmann, 1961; Kartha & Parthasarathy, 1965; North, 1965; Matthews, 1966; Hendrickson, 1979) and, as in the case of the isomorphous replacement method, the most useful applications of anomalous scattering during the last two decades have been perhaps in the field of macromolecular crystallography (Kartha, 1975; Watenpaugh, Sieker & Jensen, 1975; Vijayan, 1981). In addition to anomalous scattering of X-rays, that of neutrons was also found to have interesting applications (Koetle & Hamilton, 1975; Sikka & Rajagopal, 1975). More recently there has been a further revival in the development of anomalous-scattering methods with the advent of synchrotron radiation, particularly in view of the possibility of choosing any desired wavelength from a synchrotron-radiation source (Helliwell, 1984).

It is clear from the foregoing that the isomorphous replacement and the anomalous-scattering methods have a long and distinguished history. It is therefore impossible to do full justice to them in a comparatively short presentation like the present one. Several procedures for the application of these methods have been developed at different times. Many, although of considerable historical importance, are not extensively used at present for a variety of reasons. No attempt has been made to discuss them in detail here; the emphasis is primarily on the state of the art as it exists now. The available literature on isomorphous replacement and anomalous scattering is extensive. The reference list given at the end of the article is representative rather than exhaustive.

2.4.2. Isomorphous replacement method

2.4.2.1. Isomorphous replacement and isomorphous addition

Two crystals are said to be isomorphous if (a) both have the same space group and unit-cell dimensions and (b) the types and the positions of atoms in both are the same except for a replacement of one or more atoms in one structure with different types of atoms in the other (isomorphous replacement) or the presence of one or more additional atoms in one of them (isomorphous addition). Consider two crystal structures with identical space groups and unit-cell dimensions, one containing N atoms and the other M atoms. The N atoms in the first structure contain subsets P and Q whereas the M atoms in the second structure contain subsets P' , Q' , and R . The subset P is common to both structures in terms of atomic positions and atom types. The atomic positions are identical in subsets Q and Q' , but at any given atomic position the atom type is different in Q and Q' . The subset R exists only in the second structure. If F_N and F_M denote the structure factors of the two structures for a given reflection,

$$F_N = F_P + F_Q \quad (2.4.2.1)$$

and

$$F_M = F_P + F_{Q'} + F_R \quad (2.4.2.2)$$

2.4. ISOMORPHOUS REPLACEMENT AND ANOMALOUS SCATTERING




Fig. 2.4.2.1. Vector relationship between F_N and $F_M (= F_{NH})$.

where the quantities on the right-hand side represent contributions from different subsets. From (2.4.2.1) and (2.4.2.2) we have

$$F_M - F_N = F_H = F_{Q'} - F_Q + F_R. \quad (2.4.2.3)$$

The above equations are illustrated in the Argand diagram shown in Fig. 2.4.2.1. F_Q and $F_{Q'}$ would be collinear if all the atoms in Q were of the same type and those in Q' of another single type, as in the replacement of chlorine atoms in a structure by bromine atoms.

We have a case of 'isomorphous replacement' if $F_R = 0$ ($F_H = F_{Q'} - F_Q$) and a case of 'isomorphous addition' if $F_Q = F_{Q'} = 0$ ($F_H = F_R$). Once F_H is known, in addition to the magnitudes of F_N and F_M , which can be obtained experimentally, the two cases can be treated in an equivalent manner in reciprocal space. In deference to common practice, the term 'isomorphous replacement' will be used to cover both cases. Also, in as much as F_M is the vector sum of F_N and F_H , F_M and F_{NH} will be used synonymously. Thus

$$F_M = F_{NH} = F_N + F_H. \quad (2.4.2.4)$$

2.4.2.2. Single isomorphous replacement method

The number of replaceable (or 'added') atoms is usually small and they generally have high atomic numbers. Their positions are often determined by a Patterson synthesis of one type or another (see Chapter 2.3). It will therefore be assumed in the following discussion that F_H is known. Then it can be readily seen by referring to Fig. 2.4.2.2 that

$$\alpha_N = \alpha_H - \cos^{-1} \frac{F_{NH}^2 - F_N^2 - F_H^2}{2F_N F_H} = \alpha_H \pm \varphi; \quad (2.4.2.5)$$

when φ is derived from its cosine function, it could obviously be positive or negative. Hence, there are two possible solutions for α_N . These two solutions are distributed symmetrically about F_H . One of these would correspond to the correct value of α_N . Therefore, in general, the phase angle cannot be unambiguously determined using a pair of isomorphous crystals.

The twofold ambiguity in phase angle vanishes when the structures are centrosymmetric. F_{NH} , F_N , and F_H are all real in centric data and the corresponding phase angles are 0 or 180°. From (2.4.2.4), we then have

$$F_{NH} \pm F_N = F_H. \quad (2.4.2.6)$$




Fig. 2.4.2.2. Relationship between α_N , α_H , and φ .

The sign of F_H is already known and the signs of F_{NH} and F_N can be readily determined from (2.4.2.6) (Robertson & Woodward, 1937).

When the data are acentric, the best one can do is to use both the possible phase angles simultaneously in a Fourier synthesis (Bokhoven *et al.*, 1951). This double-phased synthesis, which is equivalent to the isomorphous synthesis of Ramachandran & Raman (1959), contains the structure and its inverse when the replaceable atoms have a centrosymmetric distribution (Ramachandran & Srinivasan, 1970). When the distribution is noncentrosymmetric, however, the synthesis contains peaks corresponding to the structure and general background. Fourier syntheses computed using the single isomorphous replacement method of Blow & Rossmann (1961) and Kartha (1961) have the same properties. In this method, the phase angle is taken to be the average of the two possible solutions of α_N , which is always α_H or $\alpha_H + 180^\circ$. Also, the Fourier coefficients are multiplied by $\cos \varphi$, following arguments based on the Blow & Crick (1959) formulation of phase evaluation (see Subsection 2.4.4.4). Although Blow & Rossmann (1961) have shown that this method could yield interpretable protein Fourier maps, it is rarely used as such in protein crystallography as the Fourier maps computed using it usually have unacceptable background levels (Blundell & Johnson, 1976).

2.4.2.3. Multiple isomorphous replacement method

The ambiguity in α_N in a noncentrosymmetric crystal can be resolved only if at least two crystals isomorphous to it are available (Bokhoven *et al.*, 1951). We then have two equations of the type (2.4.2.5), namely,

$$\alpha_N = \alpha_{H1} \pm \varphi_1 \quad \text{and} \quad (2.4.2.7)$$

$$\alpha_N = \alpha_{H2} \pm \varphi_2,$$

where subscripts 1 and 2 refer to isomorphous crystals 1 and 2, respectively. This is demonstrated graphically in Fig. 2.4.2.3 with the aid of the Harker (1956) construction. A circle is drawn with F_N as radius and the origin of the vector diagram as the centre. Two more circles are drawn with F_{NH1} and F_{NH2} as radii and the ends of vectors $-F_{H1}$ and $-F_{H2}$, respectively as centres. Each of these circles intersects the F_N circle at two points corresponding to the two possible solutions. One of the points of intersection is common and this point defines the correct value of α_N . With the assumption of perfect

2. RECIPROCAL SPACE IN CRYSTAL-STRUCTURE DETERMINATION




Fig. 2.4.2.3. Harker construction when two heavy-atom derivatives are available.

isomorphism and if errors are neglected, the phase circles corresponding to all the crystals would intersect at a common point if a number of isomorphous crystals were used for phase determination.

2.4.3. Anomalous-scattering method

2.4.3.1. Dispersion correction

Atomic scattering factors are normally calculated on the assumption that the binding energy of the electrons in an atom is negligible compared to the energy of the incident X-rays and the distribution of electrons is spherically symmetric. The transition frequencies within the atom are then negligibly small compared to the frequency of the radiation used and the scattering power of each electron in the atom is close to that of a free electron. When this assumption is valid, the atomic scattering factor is a real positive number and its value decreases as the scattering angle increases because of the finite size of the atom. When the binding energy of the electrons is appreciable, the atomic scattering factor at any given angle is given by

$$f_0 + f' + if'' \quad (2.4.3.1)$$

where f_0 is a real positive number and corresponds to the atomic scattering factor for a spherically symmetric collection of free electrons in the atom. The second and third terms are, respectively, referred to as the real and the imaginary components of the 'dispersion correction' (IT IV, 1974). f' is usually negative whereas f'' is positive. For any given atom, f'' is obviously 90° ahead of the real part of the scattering factor given by

$$f = f_0 + f' \quad (2.4.3.2)$$

The variation of f' and f'' as a function of atomic number for two typical radiations is given in Fig. 2.4.3.1 (Srinivasan, 1972; Cromer, 1965). The dispersion effects are pronounced when an absorption edge of the atom concerned is in the neighbourhood of the wavelength of the incident radiation. Atoms with high atomic numbers have several absorption edges and the dispersion-correction terms in their scattering factors always have appreciable values. The values of f' and f'' do not vary appreciably with the angle of scattering as they are caused by core electrons confined to a very small volume





Fig. 2.4.3.1. Variation of (a) f' and (b) f'' as a function of atomic number for Cu $K\alpha$ and Mo $K\alpha$ radiations. Adapted from Fig. 3 of Srinivasan (1972).

around the nucleus. An atom is usually referred to as an anomalous scatterer if the dispersion-correction terms in its scattering factor have appreciable values. The effects on the structure factors or intensities of Bragg reflections resulting from dispersion corrections are referred to as anomalous-dispersion effects or anomalous-scattering effects.

2.4.3.2. Violation of Friedel's law

Consider a structure containing N atoms of which P are normal atoms and the remaining Q anomalous scatterers. Let F_P denote the contribution of the P atoms to the structure, and F_Q and F_Q^z the real and imaginary components of the contribution of the Q atoms. The relation between the different contributions to a reflection \mathbf{h} and its Friedel equivalent $-\mathbf{h}$ is illustrated in Fig. 2.4.3.2. For simplicity we assume here that all Q atoms are of the same type. The phase angle of F_Q^z is then exactly 90° ahead of that of F_Q . The structure factors of \mathbf{h} and $-\mathbf{h}$ are denoted in the figure by $F_N(+)$ and $F_N(-)$, respectively. In the absence of anomalous scattering, or when the imaginary component of the dispersion correction is zero, the magnitudes of the two structure factors are equal and

2.4. ISOMORPHOUS REPLACEMENT AND ANOMALOUS SCATTERING




Fig. 2.4.3.2. Vector diagram illustrating the violation of Friedel's law when $F_Q^z \neq 0$.

Friedel's law is obeyed; the phase angles have equal magnitudes, but opposite signs. As can be seen from Fig. 2.4.3.2, this is no longer true when F_Q^z has a non-zero value. Friedel's law is then violated. A composite view of the vector relationship for h and $-h$ can be obtained, as in Fig. 2.4.3.3, by reflecting the vectors corresponding to $-h$ about the real axis of the vector diagram. F_P and F_Q corresponding to the two reflections superpose exactly, but F_Q^z do not. $F_N(+)$ and $F_N(-)$ then have different magnitudes and phases.

It is easily seen that Friedel's law is obeyed in centric data even when anomalous scatterers are present. F_P and F_Q are then parallel to the real axis and F_Q^z perpendicular to it. The vector sum of the three components is the same for h and $-h$. It may, however, be noted that the phase angle of the structure factor is then no longer 0 or 180°. Even when the structure is noncentrosymmetric, the effect of anomalous scattering in terms of intensity differences between Friedel equivalents varies from reflection to reflection. The difference between $F_N(+)$ and $F_N(-)$ is zero when $\alpha_P = \alpha_Q$ or $\alpha_Q + 180^\circ$. The




Fig. 2.4.3.3. A composite view of the vector relationship between $F_N(+)$ and $F_N(-)$.

difference tends to the maximum possible value ($2F_Q^z$) when $\alpha_P = \alpha_Q \pm 90^\circ$.

Intensity differences between Friedel equivalents depend also on the ratio (in terms of number and scattering power) between anomalous and normal scatterers. Differences obviously do not occur when all the atoms are normal scatterers. On the other hand, a structure containing only anomalous scatterers of the same type also does not give rise to intensity differences. Expressions for intensity differences between Friedel equivalents have been derived by Zachariasen (1965) for the most general case of a structure containing normal as well as different types of anomalous scatterers. Statistical distributions of such differences under various conditions have also been derived (Parthasarathy & Srinivasan, 1964; Parthasarathy, 1967). It turns out that, with a single type of anomalous scatterer in the structure, the ratio

$$\frac{|F_N^z(+)-F_N^z(-)|}{F_N^z(+)+F_N^z(-)}$$

has a maximum mean value when the scattering powers of the anomalous scatterers and the normal scatterers are nearly the same (Srinivasan, 1972). Also, for a given ratio between the scattering powers, the smaller the number of anomalous scatterers, the higher is the mean ratio.

2.4.3.3. Friedel and Bijvoet pairs

The discussion so far has been concerned essentially with crystals belonging to space groups $P1$ and $P\bar{1}$. In the centrosymmetric space group, the crystal and the diffraction pattern have the same symmetry, namely, an inversion centre. In $P1$, however, the crystals are noncentrosymmetric while the diffraction pattern has an inversion centre, in the absence of anomalous scattering. When anomalous scatterers are present in the structure ($F_Q^z \neq 0$), Friedel's law breaks down and the diffraction pattern no longer has an inversion centre. Thus the diffraction pattern displays the same symmetry as that of the crystal in the presence of anomalous scattering. The same is true with higher-symmetry space groups also. For example, consider a crystal with space group $P222$, containing anomalous scatterers. The magnitudes of F_P are the same for all equivalent reflections; so are those of F_Q and F_Q^z . Their phase angles, however, differ from one equivalent to another, as can be seen from Table 2.4.3.1. When $F_Q^z = 0$, the magnitudes of the vector sum of F_P and F_Q are the same for all the equivalent reflections. The intensity pattern thus has point-group symmetry $2/m\ 2/m\ 2/m$. When $F_Q^z \neq 0$, the equivalent reflections can be grouped into two sets in terms of their intensities: $hkl, h\bar{k}l, h\bar{k}l, \text{ and } hkl$; and $hkl, hkl, h\bar{k}l, \text{ and } h\bar{k}l$. The equivalents belonging to the first group have the same intensity; so have the equivalents in the second group. But the two intensities are different. Thus the symmetry of the pattern is 222 , the same as that of the crystal.

Table 2.4.3.1. Phase angles of different components of the structure factor in space group $P222$

Reflection	Phase angle ($^\circ$) of		
	F_P	F_Q	F_Q^z
$hkl, h\bar{k}l$ $h\bar{k}l, hkl$	α_P	α_Q	$90 + \alpha_Q$
$h\bar{k}l, hkl$ $hkl, h\bar{k}l$	$-\alpha_P$	$-\alpha_Q$	$90 - \alpha_Q$

2. RECIPROCAL SPACE IN CRYSTAL-STRUCTURE DETERMINATION

In general, under conditions of anomalous scattering, equivalent reflections generated by the symmetry elements in the crystal have intensities different from those of equivalent reflections generated by the introduction of an additional inversion centre in normal scattering. There have been suggestions that a reflection from the first group and another from the second group should be referred to as a 'Bijvoet pair' instead of a 'Friedel pair', when the two reflections are not inversely related. Most often, however, the terms are used synonymously. The same practice will be followed in this article.

2.4.3.4. Determination of absolute configuration

The determination of the absolute configuration of chiral molecules has been among the most important applications of anomalous scattering. Indeed, anomalous scattering is the only effective method for this purpose and the method, first used in the early 1950's (Peerdeman, van Bommel & Bijvoet, 1951), has been extensively employed in structural crystallography (Ramaseshan, 1963; Vos, 1975).

Many molecules, particularly biologically important ones, are chiral in that the molecular structure is not superimposable on its mirror image. Chirality (handedness) arises primarily on account of the presence of asymmetric carbon atoms in the molecule. A tetravalent carbon is asymmetric when the four atoms (or groups) bonded to it are all different from one another. The substituents can then have two distinct arrangements which are mirror images of (or related by inversion to) each other. These optical isomers or enantiomers have the same chemical and physical properties except that they rotate the plane of polarization in opposite directions when polarized light passes through them. It is not, however, possible to calculate the sign of optical rotation, given the exact spatial arrangement or the 'absolute configuration' of the molecule. Therefore, one cannot distinguish between the possible enantiomeric configurations of a given asymmetric molecule from measurements of optical rotation. This is also true of molecules with chiralities generated by overall asymmetric geometry instead of the presence of asymmetric carbon atoms in them.

Normal X-ray scattering does not distinguish between enantiomers. Two structures $A(x_j, y_j, z_j)$ and $B(-x_j, -y_j, -z_j)$ ($j=1 \dots N$) obviously produce the same diffraction pattern on account of Friedel's law. The situation is, however, different when anomalous scatterers are present in the structure. The intensity difference between reflections h and $-h$, or that between members of any Bijvoet pair, has the same magnitude, but opposite sign for structures A and B . If the atomic coordinates are known, the intensities of Bijvoet pairs can be readily calculated. The absolute configuration can then be determined, i.e. one can distinguish between A and B by comparing the calculated intensities with the observed ones, for a few Bijvoet pairs with pronounced anomalous effects.

2.4.3.5. Determination of phase angles

An important application of anomalous scattering is in the determination of phase angles using Bijvoet differences (Ramachandran & Raman, 1956; Peerdeman & Bijvoet, 1956). From Figs. 2.4.3.2 and 2.4.3.3, we have

$$F_N^2(+) = F_N^2 + F_Q^2 + 2F_N F_Q'' \cos \theta \quad (2.4.3.3)$$

and

$$F_N^2(-) = F_N^2 + F_Q^2 - 2F_N F_Q'' \cos \theta. \quad (2.4.3.4)$$

Then

$$\cos \theta = \frac{F_N^2(+) - F_N^2(-)}{4F_N F_Q''}. \quad (2.4.3.5)$$

In the above equations F_N may be approximated to $[F_N(+) + F_N(-)]/2$. Then θ can be evaluated from (2.4.3.5) except for the ambiguity in its sign. Therefore, from Fig. 2.4.3.2, we have

$$\alpha_N = \alpha_Q + 90^\circ \pm \theta. \quad (2.4.3.6)$$

The phase angle thus has two possible values symmetrically distributed about F_Q'' . Anomalous scatterers are usually heavy atoms and their positions can most often be determined by Patterson methods. α_Q can then be calculated and the two possible values of α_N for each reflection evaluated using (2.4.3.6).

In practice, the twofold ambiguity in phase angles can often be resolved in a relatively straightforward manner. As indicated earlier, anomalous scatterers usually have relatively high atomic numbers. The 'heavy-atom' phases calculated from their positions therefore contain useful information. For any given reflection, that phase angle which is closer to the heavy-atom phase, from the two phases calculated using (2.4.3.6), may be taken as the correct phase angle. This method has been successfully used in several structure determinations including that of a derivative of vitamin B₁₂ (Dale *et al.*, 1963). The same method was also employed in a probabilistic fashion in the structure solution of a small protein (Hendrickson & Teeter, 1981). A method for obtaining a unique, but approximate, solution for phase angles from (2.4.3.6) has also been suggested (Srinivasan & Chacko, 1970). An accurate unique solution for phase angles can be obtained if one collects two sets of intensity data using two different wavelengths which have different dispersion-correction terms for the anomalous scatterers in the structure. Two equations of the type (2.4.3.6) are then available for each reflection and the solution common to both is obviously the correct phase angle. Different types of Patterson and Fourier syntheses can also be employed for structure solution using intensity differences between Bijvoet equivalents (Srinivasan, 1972; Okaya & Pepinsky, 1956; Pepinsky, Okaya & Takeuchi, 1957).

An interesting situation occurs when the replaceable atoms in a pair of isomorphous structures are anomalous scatterers. The phase angles can then be uniquely determined by combining isomorphous replacement and anomalous-scattering methods. Such situations normally occur in protein crystallography and are discussed in Subsection 2.4.4.5.

2.4.3.6. Anomalous scattering without phase change

The phase determination, and hence the structure solution, outlined above relies on the imaginary component of the dispersion correction. Variation in the real component can also be used in structure analysis. In early applications of anomalous scattering, the real component of the dispersion correction was made use of to distinguish between atoms of nearly the same atomic numbers (Mark & Szillard, 1925; Bradley & Rodgers, 1934). For example, copper and manganese, with atomic numbers 29 and 25, respectively, are not easily distinguishable under normal X-ray scattering. However, the real components of the dispersion correction for the two elements are -1.129 and -3.367 , respectively, when Fe K α radiation is used (IT IV, 1974). Therefore, the difference between the scattering factors of the two elements is accentuated when this radiation is used. The difference is more pronounced at high angles as the normal scattering

2.4. ISOMORPHOUS REPLACEMENT AND ANOMALOUS SCATTERING

factor falls off comparatively rapidly with increasing scattering angle whereas the dispersion-correction term does not.

The structure determination of KMnO_4 provides a typical example for the use of anomalous scattering without phase change in the determination of a centrosymmetric structure (Ramaseshan, Venkatesan & Mani, 1957; Ramaseshan & Venkatesan, 1957). f' and f'' for manganese for $\text{Cu } K\alpha$ radiation are -0.568 and 2.808 , respectively. The corresponding values for $\text{Fe } K\alpha$ radiation are -3.367 and 0.481 , respectively (IT IV, 1974). The data sets collected using the two radiations can now be treated as those arising from two perfectly isomorphous crystals. The intensity differences between a reflection in one set and the corresponding reflection in the other are obviously caused by the differences in the dispersion-correction terms. They can, however, be considered formally as intensity differences involving data from two perfectly isomorphous crystals. They can be used, as indeed they were, to determine the position of the manganese ion through an appropriate Patterson synthesis (see Subsection 2.4.4.2) and then to evaluate the signs of structure factors using (2.4.2.6) when the structure is centrosymmetric. When the structure is noncentrosymmetric, a twofold ambiguity exists in the phase angles in a manner analogous to that in the isomorphous replacement method. This ambiguity can be removed if the structure contains two different subsets of atoms Q_1 and Q_2 which, respectively, scatter radiations λ_{Q_1} and λ_{Q_2} anomalously. Data sets can then be collected with λ , which is scattered normally by all atoms, λ_{Q_1} and λ_{Q_2} . The three sets can be formally treated as those from three perfectly isomorphous structures and the phase determination effected using (2.4.2.7) (Ramaseshan, 1963).

2.4.3.7. Treatment of anomalous scattering in structure refinement

The effect of anomalous scattering needs to be taken into account in the refinement of structures containing anomalous scatterers, if accurate atomic parameters are required. The effect of the real part of the dispersion correction is largely confined to the thermal parameters of anomalous scatterers. This effect can be eliminated by simply adding f' to the normal scattering factor of the anomalous scatterers.

The effects of the imaginary component of the dispersion correction are, however, more complex. These effects could lead to serious errors in positional parameters when the space group is polar, if data in the entire diffraction sphere are not used (Ueki, Zalkin & Templeton, 1966; Cruickshank & McDonald, 1967). For example, accessible data in a hemisphere are normally used for X-ray analysis when the space group is $P1$. If the hemisphere has say h positive, the x coordinates of all the atoms would be in error when the structure contains anomalous scatterers. The situation in other polar space groups has been discussed by Cruickshank & McDonald (1967). In general, in the presence of anomalous scattering, it is desirable to collect data for the complete sphere, if accurate structural parameters are required (Srinivasan, 1972).

Methods have been derived to correct for dispersion effects in observed data from centrosymmetric and noncentrosymmetric crystals (Patterson, 1963). The methods are empirical and depend upon the refined parameters at the stage at which corrections are applied. This is obviously an unsatisfactory situation and it has been suggested that the measured structure factors of Bijvoet equivalents should instead be treated as independent observations in structure refinement (Ibers & Hamilton, 1964). The effect of dispersion corrections needs

to be taken into account to arrive at the correct scale and temperature factors also (Wilson, 1975; Gilli & Cruickshank, 1973).

2.4.4. Isomorphous replacement and anomalous scattering in protein crystallography

2.4.4.1. Protein heavy-atom derivatives

Perhaps the most spectacular applications of isomorphous replacement and anomalous-scattering methods have been in the structure solution of large biological macromolecules, primarily proteins. Since its first successful application on myoglobin and haemoglobin, the isomorphous replacement method, which is often used in conjunction with the anomalous-scattering method, has been employed in the solution of scores of proteins. The application of this method involves the preparation of protein heavy-atom derivatives, i.e. the attachment of heavy atoms like mercury, uranium, and lead or chemical groups containing them, to protein crystals in a coherent manner without changing the conformation of the molecules and their crystal packing. This is only rarely possible in ordinary crystals as the molecules in them are closely packed. Protein crystals, however, contain large solvent regions and isomorphous derivatives can be prepared by replacing the disordered solvent molecules by heavy-atom-containing groups without disturbing the original arrangement of protein molecules.

2.4.4.2. Determination of heavy-atom parameters

For any given reflection, the structure factor of the native protein crystal (F_N), that of a heavy-atom derivative (F_{NH}), and the contribution of the heavy atoms in that derivative (F_H) are related by the equation

$$F_{NH} = F_N + F_H. \quad (2.4.4.1)$$

The value of F_H depends not only on the positional and thermal parameters of the heavy atoms, but also on their occupancy factors, because, at a given position, the heavy atom may not often be present in all the unit cells. For example, if the heavy atom is present at a given position in only half the unit cells in the crystal, then the occupancy factor of the site is said to be 0.5.

For the successful determination of the heavy-atom parameters, as also for the subsequent phase determination, the data sets from the native and the derivative crystals should have the same relative scale. The different data sets should also have the same overall temperature factor. Different scaling procedures have been suggested (Blundell & Johnson, 1976) and, among them, the following procedure, based on Wilson's (1942) statistics, appears to be the most feasible in the early stages of structure analysis.

Assuming that the data from the native and the derivative crystals obey Wilson's statistics, we have, for any range of $\sin^2 \theta / \lambda^2$,

$$\ln \left\{ \frac{\sum f_{Nj}^2}{\langle F_N^2 \rangle} \right\} = \ln K_N + 2B_N \frac{\sin^2 \theta}{\lambda^2} \quad (2.4.4.2)$$

and

$$\ln \left\{ \frac{\sum f_{Nj}^2 + \sum f_{Hj}^2}{\langle F_{NH}^2 \rangle} \right\} = \ln K_{NH} + 2B_{NH} \frac{\sin^2 \theta}{\lambda^2}, \quad (2.4.4.3)$$

where f_{Nj} and f_{Hj} refer to the atomic scattering factors of protein atoms and heavy atoms, respectively. K_N and K_{NH} are the scale factors to be applied to the intensities from the

2. RECIPROCAL SPACE IN CRYSTAL-STRUCTURE DETERMINATION

native and the derivative crystals, respectively, and B_N and B_{NH} the temperature factors of the respective structure factors. Normally one would be able to derive the absolute scale factor and the temperature factor for both the data sets from (2.4.4.2) and (2.4.4.3) using the well known Wilson plot. The data from protein crystals, however, do not follow Wilson's statistics as protein molecules contain highly non-random features. Therefore, in practice, it is difficult to fit a straight line through the points in a Wilson plot, thus rendering the parameters derived from it unreliable. (2.4.4.2) and (2.4.4.3) can, however, be used in a different way. From the two equations we obtain

$$\ln \left\{ \frac{\sum f_{NH}^2 + \sum f_{H'}^2}{\sum f_{NH}^2} \cdot \frac{\langle F_N^2 \rangle}{\langle F_{NH}^2 \rangle} \right\} = \ln \left(\frac{K_{NH}}{K_N} \right) + 2(B_{NH} - B_N) \frac{\sin^2 \theta}{\lambda^2}. \quad (2.4.4.4)$$

The effects of structural non-randomness in the crystals obviously cancel out in (2.4.4.4). When the left-hand side of (2.4.4.4) is plotted against $(\sin^2 \theta)/\lambda^2$, it is called a comparison or difference Wilson plot. Such plots yield the ratio between the scales of the derivative and the native data, and the additional temperature factor of the derivative data. Initially, the number and the occupancy factors of heavy-atom sites are unknown, and are roughly estimated from intensity differences to evaluate $\sum f_{H'}^2$. These estimates usually undergo considerable revision in the course of the determination and the refinement of heavy-atom parameters.

At first, heavy-atom positions are most often determined by Patterson syntheses of one type or another. Such syntheses are discussed in some detail elsewhere in Chapter 2.3. They are therefore discussed here only briefly.

Equation (2.4.2.6) holds when the data are centric. F_H is usually small compared to F_N and F_{NH} , and the minus sign is then relevant on the left-hand side of (2.4.2.6). Thus the difference between the magnitudes of F_{NH} and F_N , which can be obtained experimentally, normally gives a correct estimate of the magnitude of F_H for most reflections. Then a Patterson synthesis with $(F_{NH} - F_N)^2$ as coefficients corresponds to the distribution of vectors between heavy atoms, when the data are centric. But proteins are made up of L-amino acids and hence cannot crystallize in centrosymmetric space groups. However, many proteins crystallize in space groups with centrosymmetric projections. The centric data corresponding to these projections can then be used for determining heavy-atom positions through a Patterson synthesis of the type outlined above.

The situation is more complex for three-dimensional acentric data. It has been shown (Rossmann, 1961) that

$$(F_{NH} - F_N)^2 = F_H^2 \cos^2(\alpha_{NH} - \alpha_H) \quad (2.4.4.5)$$

when F_H is small compared to F_{NH} and F_N . Patterson synthesis with $(F_{NH} - F_N)^2$ as coefficients would, therefore, give an approximation to the heavy-atom vector distribution. An isomorphous difference Patterson synthesis of this type has been used extensively in protein crystallography to determine heavy-atom positions. The properties of this synthesis have been extensively studied (Ramachandran & Srinivasan, 1970; Rossmann, 1960; Phillips, 1966; Dodson & Vijayan, 1971) and it has been shown that this Patterson synthesis would provide a good approximation to the heavy-atom vector distribution even when F_H is large compared to F_N (Dodson & Vijayan, 1971).

As indicated earlier (see Subsection 2.4.3.1), heavy atoms are always anomalous scatterers, and the structure factors of

any given reflection and its Friedel equivalent from a heavy-atom derivative have unequal magnitudes. If these structure factors are denoted by $F_{NH}(+)$ and $F_{NH}(-)$ and the real component of the heavy-atom contributions (including the real component of the dispersion correction) by F_H , then it can be shown (Kartha & Parthasarathy, 1965) that

$$\left(\frac{k}{2}\right)^2 [(F_{NH}(+) - F_{NH}(-))]^2 = F_H^2 \sin^2(\alpha_{NH} - \alpha_H), \quad (2.4.4.6)$$

where $k = (f_H + f'_H)/f'_H$. Here it has been assumed that all the anomalous scatterers are of the same type with atomic scattering factor f_H and dispersion correction terms f'_H and f''_H . A Patterson synthesis with the left-hand side of (2.4.4.6) as coefficients would also yield the vector distribution corresponding to the heavy-atom positions (Rossmann, 1961; Kartha & Parthasarathy, 1965). However, $F_{NH}(+) - F_{NH}(-)$ is a small difference between two large quantities and is liable to be in considerable error. Patterson syntheses of this type are therefore rarely used to determine heavy-atom positions.

It is interesting to note (Kartha & Parthasarathy, 1965) that addition of (2.4.4.5) and (2.4.4.6) readily leads to

$$(F_{NH} - F_N)^2 + \left(\frac{k}{2}\right)^2 [F_{NH}(+) - F_{NH}(-)]^2 = F_H^2. \quad (2.4.4.7)$$

Thus, the magnitude of the heavy-atom contribution can be estimated if intensities of Friedel equivalents have been measured from the derivative crystal. F_{NH} is then not readily available, but to a good approximation

$$F_{NH} = [F_{NH}(+) + F_{NH}(-)]/2. \quad (2.4.4.8)$$

A different and more accurate expression for estimating F_H^2 from isomorphous and anomalous differences was derived by Matthews (1966). According to a still more accurate expression derived by Singh & Ramaseshan (1966),

$$\begin{aligned} F_H^2 &= F_{NH}^2 + F_N^2 - 2F_{NH}F_N \cos(\alpha_N - \alpha_{NH}) \\ &= F_{NH}^2 + F_N^2 \pm 2F_{NH}F_N \\ &\quad \times (1 - \{k[F_{NH}(+) - F_{NH}(-)]/2F_N\}^2)^{1/2}. \end{aligned} \quad (2.4.4.9)$$

The lower estimate in (2.4.4.9) is relevant when $|\alpha_N - \alpha_{NH}| < 90^\circ$ and the upper estimate is relevant when $|\alpha_N - \alpha_{NH}| > 90^\circ$. The lower and the upper estimates may be referred to as F_{HLE} and F_{HUE} , respectively. It can be readily shown (Dodson & Vijayan, 1971) that the lower estimate would represent the correct value of F_H for a vast majority of reflections. Thus, a Patterson synthesis with F_{HLE}^2 as coefficients would yield the vector distribution of heavy atoms in the derivative. Such a synthesis would normally be superior to those with the left-hand sides of (2.4.4.5) and (2.4.4.6) as coefficients. However, when the level of heavy-atom substitution is low, the anomalous differences are also low and susceptible to large percentage errors. In such a situation, a synthesis with $(F_{NH} - F_N)^2$ as coefficients is likely to yield better results than that with F_{HLE}^2 as coefficients (Vijayan, 1981).

Direct methods employing different methodologies have also been used successfully for the determination of heavy-atom positions (Navia & Sigler, 1974). These methods, developed primarily for the analysis of smaller structures, have not yet been successful in *a priori* analysis of protein structures. The very size of protein structures makes the probability relations used in these methods weak. In addition, data from protein crystals do not normally extend to high enough angles to permit resolution of individual atoms in the structure and the feasibility of using many of the currently popular

2.4. ISOMORPHOUS REPLACEMENT AND ANOMALOUS SCATTERING

direct-method procedures in such a situation has been a topic of much discussion. The heavy atoms in protein derivative crystals, however, are small in number and are normally situated far apart from one another. They are thus expected to be resolved even when low-resolution X-ray data are used. In most applications, the magnitudes of the differences between F_{NH} and F_N are formally considered as the 'observed structure factors' of the heavy-atom distribution and conventional direct-method procedures are then applied to them.

Once the heavy-atom parameters in one or more derivatives have been determined, approximate protein phase angles, α_N 's, can be derived using methods described later. These phase angles can then be readily used to determine the heavy-atom parameters in a new derivative employing a difference Fourier synthesis with coefficients

$$(F_{NH} - F_N) \exp(i\alpha_N). \quad (2.4.4.10)$$

Such syntheses are also used to confirm and to improve upon the information on heavy-atom parameters obtained through Patterson or direct methods. They are obviously very powerful when centric data corresponding to centrosymmetric projections are used. The synthesis yields satisfactory results even when the data are acentric although the difference Fourier technique becomes progressively less powerful as the level of heavy-atom substitution increases (Dodson & Vijayan, 1971).

While the positional parameters of heavy atoms can be determined with a reasonable degree of confidence using the above-mentioned methods, the corresponding temperature and occupancy factors cannot. Rough estimates of the latter are usually made from the strength and the size of appropriate peaks in difference syntheses. The estimated values are then refined, along with the positional parameters, using the techniques outlined below.

2.4.4.3. Refinement of heavy-atom parameters

The least-squares method with different types of minimization functions is used for refining the heavy-atom parameters, including the occupancy factors. The most widely used method (Dickerson, Kendrew & Strandberg, 1961; Muirhead, Cox, Mazzarella & Perutz, 1967; Dickerson, Weinzierl & Palmer, 1968) involves the minimization of the function

$$\varphi = \sum w(F_{NH} - |F_N + F_H|)^2, \quad (2.4.4.11)$$

where the summation is over all the reflections and w is the weight factor associated with each reflection. Here F_{NH} is the observed magnitude of the structure factor for the particular derivative and $F_N + F_H$ is the calculated structure factor. The latter obviously depends upon the protein phase angle α_N , and the magnitude and the phase angle of F_H which are in turn dependent on the heavy-atom parameters. Let us assume that we have three derivatives A , B , and C , and that we have already determined the heavy-atom parameters HA_i , HB_i , and HC_i . Then,

$$\begin{aligned} F_{HA} &= F_{HA}(HA_i) \\ F_{HB} &= F_{HB}(HB_i) \\ F_{HC} &= F_{HC}(HC_i). \end{aligned} \quad (2.4.4.12)$$

A set of approximate protein phase angles is first calculated, employing methods described later, making use of the unrefined heavy-atom parameters. These phase angles are used to construct $F_N + F_H$ for each derivative. (2.4.4.11) is then minimized, separately for each derivative, by varying HA_i for derivative A , HB_i for derivative B , and HC_i for derivative C . The refined values of HA_i , HB_i , and HC_i are

subsequently used to calculate a new set of protein phase angles. Alternate cycles of parameter refinement and phase-angle calculation are carried out until convergence is reached. The progress of refinement may be monitored by computing an R factor defined as (Kraut, Sieker, High & Freer, 1962)

$$R_K = \frac{\sum |F_{NH} - |F_N + F_H||}{F_{NH}} \quad (2.4.4.13)$$

The above method has been successfully used for the refinement of heavy-atom parameters in the X-ray analysis of many proteins. However, it has one major drawback in that the refined parameters in one derivative are dependent on those in other derivatives through the calculation of protein phase angles. Therefore, it is important to ensure that the derivative, the heavy-atom parameters of which are being refined, is omitted from the phase-angle calculation (Blow & Matthews, 1973). Even when this is done, serious problems might arise when different derivatives are related by common sites. In practice, the occupancy factors of the common sites tend to be overestimated compared to those of the others (Vijayan, 1981; Dodson & Vijayan, 1971). Yet another factor which affects the occupancy factors is the accuracy of the phase angles. The inclusion of poorly phased reflections tends to result in the underestimation of occupancy factors. It is therefore advisable to omit from refinement cycles reflections with figures of merit less than a minimum threshold value or to assign a weight proportional to the figure of merit (as defined later) to each term in the minimization function (Dodson & Vijayan, 1971; Blow & Matthews, 1973).

If anomalous-scattering data from derivative crystals are available, the values of F_H can be estimated using (2.4.4.7) or (2.4.4.9) and these can be used as the 'observed' magnitudes of the heavy-atom contributions for the refinement of heavy-atom parameters, as has been done by many workers (Watenpaugh *et al.*, 1975; Vijayan, 1981; Kartha, 1965). If (2.4.4.9) is used for estimating F_H , the minimization function has the form

$$\varphi = \sum w(F_{HLE} - F_H)^2. \quad (2.4.4.14)$$

The progress of refinement may be monitored using a reliability index defined as

$$R = \frac{\sum |F_{HLE} - F_H|}{\sum F_{HLE}}. \quad (2.4.4.15)$$

The major advantage of using F_{HLE} 's in refinement is that the heavy-atom parameters in each derivative can now be refined independently of all other derivatives. Care should, however, be taken to omit from calculations all reflections for which F_{HLE} is likely to be the correct estimate of F_H . This can be achieved in practice by excluding from least-squares calculations all reflections for which F_{HLE} has a value less than the maximum expected value of F_H for the given derivative (Vijayan, 1981; Dodson & Vijayan, 1971).

A major problem associated with this refinement method is concerned with the effect of experimental errors on refined parameters. The values of $F_{NH}(+) - F_{NH}(-)$ are often comparable to the experimental errors associated with $F_{NH}(+)$ and $F_{NH}(-)$. In such a situation, even random errors in $F_{NH}(+)$ and $F_{NH}(-)$ tend to increase systematically the observed difference between them (Dodson & Vijayan, 1971). In (2.4.4.7) and (2.4.4.9), this difference is multiplied by k or $k/2$, a quantity much greater than unity, and then squared. This could lead to the systematic overestimation of F_{HLE} 's and the consequent overestimation of occupancy factors. The situation can be improved by employing empirical values of

2. RECIPROCAL SPACE IN CRYSTAL-STRUCTURE DETERMINATION

k , evaluated using the relation (Kartha & Parthasarathy, 1965; Mathews, 1966)

$$k = \frac{2 \sum |F_{NH} - F_N|}{\sum |F_{NH}(+) - F_{NH}(-)|}, \quad (2.4.4.16)$$

for estimating F_{HLE} or by judiciously choosing the weighting factors in (2.4.4.14) (Dodson & Vijayan, 1971). The use of a modified form of F_{HLE} , arrived at through statistical considerations, along with appropriate weighting factors, has also been advocated (Dodson, Evans & French, 1975).

When the data are centric, (2.4.4.9) reduces to

$$F_H = F_{NH} \pm F_N. \quad (2.4.4.17)$$

Here, again, the lower estimate most often corresponds to the correct value of F_H . (2.4.4.17) does not involve $F_{NH}(+) - F_{NH}(-)$ which, as indicated earlier, is prone to substantial error. Therefore, F_H 's estimated using centric data are more reliable than those estimated using acentric data. Consequently, centric reflections, when available, are extensively used for the refinement of heavy-atom parameters. It may also be noted that in conditions under which F_{HLE} corresponds to the correct estimate of F_H , minimization functions (2.4.4.11) and (2.4.4.14) are identical for centric data.

A Patterson function correlation method with a minimization function of the type

$$\varphi = \sum w[(F_{NH} - F_N)^2 - F_H^2]^2 \quad (2.4.4.18)$$

was among the earliest procedures suggested for heavy-atom parameter refinement (Rossmann, 1960). This procedure would obviously work well when centric reflections are used. A modified version of this procedure, in which the origins of the Patterson functions are removed from the correlation, and centric and acentric data are treated separately, has been proposed recently (Terwilliger & Eisenberg, 1983).

2.4.4.4. Treatment of errors in phase evaluation: Blow and Crick formulation

As shown in Subsection 2.4.2.3, ideally protein phase angles can be evaluated if two isomorphous heavy-atom derivatives are available. However, in practice, conditions are far from ideal on account of several factors such as imperfect isomorphism, errors in the estimation of heavy-atom parameters, and the experimental errors in the measurement of intensity from the native and the derivative crystals. It is therefore desirable to use as many derivatives as are available for phase determination. When isomorphism is imperfect and errors exist in data and heavy-atom parameters, none of the circles in a Harker diagram would intersect at a single point; instead, there would be a distribution of intersections, such as that illustrated in Fig. 2.4.4.1. Consequently, a unique solution for the phase angle cannot be deduced.

The statistical procedure for computing protein phase angles using multiple isomorphous replacement (MIR) was derived by Blow & Crick (1959). In their treatment, Blow and Crick assume, for mathematical convenience, that all errors, including those arising from imperfect isomorphism, could be considered as residing in the magnitudes of the derivative structure factors only. They further assume that these errors could be described by a Gaussian distribution. With these simplifying assumptions, the statistical procedure for phase determination could be derived in the following manner.

Consider the vector diagram, shown in Fig. 2.4.4.2, for a reflection from the i th derivative for an arbitrary value α for the protein phase angle. Then,

$$D_{Hi}(\alpha) = [F_N^2 + F_{Hi}^2 + 2F_N F_{Hi} \cos(\alpha_{Hi} - \alpha)]^{1/2}. \quad (2.4.4.19)$$




Fig. 2.4.4.1. Distribution of intersections in the Harker construction under non-ideal conditions.




Fig. 2.4.4.2. Vector diagram indicating the calculated structure factor, $D_{Hi}(\alpha)$, of the i th heavy-atom derivative for an arbitrary value α for the phase angle of the structure factor of the native protein.

If α corresponds to the true protein phase angle α_N , then D_{Hi} coincides with F_{NH_i} . The amount by which $D_{Hi}(\alpha)$ differs from F_{NH_i} , namely,

$$\xi_{Hi}(\alpha) = F_{NH_i} - D_{Hi}(\alpha), \quad (2.4.4.20)$$

is a measure of the departure of α from α_N , ξ is called the lack of closure. The probability for α being the correct protein phase angle could now be defined as

$$P_i(\alpha) = N_i \exp[-\xi_{Hi}^2(\alpha)/2E_i^2], \quad (2.4.4.21)$$

where N_i is the normalization constant and E_i is the estimated r.m.s. error. The methods for estimating E_i will be outlined later. When several derivatives are used for phase determination, the total probability of the phase angle α being the protein phase angle would be

$$P(\alpha) = \prod P_i(\alpha) = N \exp\left\{-\sum_i [\xi_{Hi}^2(\alpha)/2E_i^2]\right\}, \quad (2.4.4.22)$$

where the summation is over all the derivatives. A typical distribution of $P(\alpha)$ plotted around a circle of unit radius is

2.4. ISOMORPHOUS REPLACEMENT AND ANOMALOUS SCATTERING




Fig. 2.4.4.3. The probability distribution of the protein phase angle. The point P is the centroid of the distribution.

shown in Fig. 2.4.4.3. The phase angle corresponding to the highest value of $P(\alpha)$ would obviously be the most probable protein phase, α_M , of the given reflection. The most probable electron-density distribution is obtained if each F_N is associated with the corresponding α_M , in a Fourier synthesis.

Blow and Crick suggested a different way of using the probability distribution. In Fig. 2.4.4.3, the centroid of the probability distribution is denoted by P . The polar coordinates of P are m and α_B , where m , a fractional positive number with a maximum value of unity, and α_B are referred to as the 'figure of merit' and the 'best phase', respectively. One can then compute a 'best Fourier' with coefficients

$$mF_N \exp(i\alpha_B).$$

The best Fourier is expected to provide an electron-density distribution with the lowest r.m.s. error. The figure of merit and the best phase are usually calculated using the equations

$$\begin{aligned} m \cos \alpha_B &= \frac{\sum_i P(\alpha_i) \cos(\alpha_i)}{\sum_i P(\alpha_i)} \\ m \sin \alpha_B &= \frac{\sum_i P(\alpha_i) \sin(\alpha_i)}{\sum_i P(\alpha_i)}, \end{aligned} \quad (2.4.4.23)$$

where $P(\alpha_i)$ are calculated, say, at 5° intervals (Dickerson *et al.*, 1961). The figure of merit is statistically interpreted as the cosine of the expected error in the calculated phase angle and it is obviously a measure of the precision of phase determination. In general, m is high when α_M and α_B are close to each other and low when they are far apart.

2.4.4.5. Use of anomalous scattering in phase evaluation

When anomalous-scattering data have been collected from derivative crystals, $F_{NH}(+)$ and $F_{NH}(-)$ can be formally treated as arising from two independent derivatives. The corresponding Harker diagram is shown in Fig. 2.4.4.4. Thus, in principle, protein phase angles can be determined using a single derivative when anomalous-scattering effects are also made use of. It is interesting to note that the information obtained from isomorphous differences, $F_{NH} - F_N$, and that obtained from anomalous differences, $F_{NH}(+) - F_{NH}(-)$, are complementary. The isomorphous difference for any given reflection is a maximum when F_N and F_H are parallel or antiparallel. The anomalous difference is then zero, if all the anomalous scatterers are of the same type, and α_N is determined uniquely on the basis of the isomorphous difference.




Fig. 2.4.4.4. Harker construction using anomalous-scattering data from a single derivative.

The isomorphous difference decreases and the anomalous difference increases as the inclination between F_N and F_H increases. The isomorphous difference tends to be small and the anomalous difference tends to have the maximum possible value when F_N and F_H are perpendicular to each other. The anomalous difference then has the predominant influence in determining the phase angle.

Although isomorphous and anomalous differences have a complementary role in phase determination, their magnitudes are obviously unequal. Therefore, when $F_{NH}(+)$ and $F_{NH}(-)$ are treated as arising from two derivatives, the effect of anomalous differences on phase determination would be only marginal as, for any given reflection, $F_{NH}(+) - F_{NH}(-)$ is usually much smaller than $F_{NH} - F_N$. However, the magnitude of the error in the anomalous difference would normally be much smaller than that in the corresponding isomorphous difference. Firstly, the former is obviously free from the effects of imperfect isomorphism. Secondly, $F_{NH}(+)$ and $F_{NH}(-)$ are expected to have the same systematic errors as they are measured from the same crystal. These errors are eliminated in the difference between the two quantities. Therefore, as pointed out by North (1965), the r.m.s. error used for anomalous differences should be much smaller than that used for isomorphous differences. Denoting the r.m.s. error in anomalous differences by E' , the new expression for the probability distribution of protein phase angle may be written as

$$\begin{aligned} P_i(\alpha) &= N_i \exp[-\xi_{FH}^2(\alpha)/2E_i^2] \\ &\times \exp\{-[\Delta H_i - \Delta H_{ical}(\alpha)]^2/2E_i'^2\}, \end{aligned} \quad (2.4.4.24)$$

where

$$\Delta H_i = F_{NH}(+) - F_{NH}(-)$$

and

$$\Delta H_{ical}(\alpha) = 2F_H'' \sin(\alpha_{Di} - \alpha_{Hi}).$$

Here α_{Di} is the phase angle of $D_{Hi}(\alpha)$ [see (2.4.4.19) and Fig. 2.4.4.2]. $\Delta H_{ical}(\alpha)$ is the anomalous difference calculated for the assumed protein phase angle α . F_{NH} may be taken as the average of $F_{NH}(+)$ and $F_{NH}(-)$ for calculating $\xi_{FH}^2(\alpha)$ using (2.4.4.20).

2.4.4.6. Estimation of r.m.s. error

Perhaps the most important parameters that control the reliability of phase evaluation using the Blow and Crick

2. RECIPROCAL SPACE IN CRYSTAL-STRUCTURE DETERMINATION

formulations are the isomorphous r.m.s. error E_i and the anomalous r.m.s. error E'_i . For a given derivative, the sharpness of the peak in the phase probability distribution obviously depends upon the value of E and that of E' when anomalous-scattering data have also been used. When several derivatives are used, an overall underestimation of r.m.s. errors leads to artificially sharper peaks, the movement of α_B towards α_M , and deceptively high figures of merit. Opposite effects result when E 's are overestimated. Underestimation or overestimation of the r.m.s. error in the data from a particular derivative leads to distortions in the relative contribution of that derivative to the overall phase probability distributions. It is therefore important that the r.m.s. error in each derivative is correctly estimated.

Centric reflections, when present, obviously provide the best means for evaluating E using the expression

$$E^2 = \sum_n (|F_{NH} \pm F_N| - F_N)^2 / n \quad (2.4.4.25)$$

As suggested by Blow & Crick (1959), values of E thus estimated can be used for acentric reflections as well. Once a set of approximate protein phase angles is available E_i can be calculated as the r.m.s. lack of closure corresponding to α_B [i.e. $\alpha = \alpha_B$ in (2.4.4.20)] (Kartha, 1976). E'_i can be similarly evaluated as the r.m.s. difference between the observed anomalous difference and the anomalous difference calculated for α_B [see (2.4.4.24)]. Normally, the value of E'_i is about a third of that of E_i (North, 1965).

A different method, outlined below, can also be used to evaluate E and E' when anomalous scattering is present (Vijayan, 1981; Adams, 1968). From Fig. 2.4.2.2, we have

$$\cos \psi = (F_{NH}^2 + F_H^2 - F_N^2) / 2F_{NH}F_H \quad (2.4.4.26)$$

and

$$F_N^2 = F_{NH}^2 + F_H^2 - 2F_{NH}F_H \cos \psi, \quad (2.4.4.27)$$

where $\psi = \alpha_{NH} - \alpha_H$. Using arguments similar to those used in deriving (2.4.3.5), we obtain

$$\sin \psi = [F_{NH}^2(+) - F_{NH}^2(-)] / 4F_{NH}F_H'. \quad (2.4.4.28)$$

If F_{NH} is considered to be equal to $[F_{NH}(+) + F_{NH}(-)] / 2$, we obtain from (2.4.4.28)

$$F_{NH}(+) - F_{NH}(-) = 2F_H' \sin \psi. \quad (2.4.4.29)$$

We obtain what may be called ψ_{iso} if the magnitude of ψ is determined from (2.4.4.26) and the quadrant from (2.4.4.28). Similarly, we obtain ψ_{ano} if the magnitude of ψ is determined from (2.4.4.28) and the quadrant from (2.4.4.26). Ideally, ψ_{iso} and ψ_{ano} should have the same value and the difference between them is a measure of the errors in the data. F_N obtained from (2.4.4.27) using ψ_{ano} may be considered as its calculated value (F_{Ncal}). Then, assuming all errors to lie in F_N , we may write

$$E^2 = \sum_n (F_N - F_{Ncal})^2 / n. \quad (2.4.4.30)$$

Similarly, the calculated anomalous difference (ΔH_{cal}) may be evaluated from (2.4.4.29) using ψ_{iso} . Then

$$E^2 = \sum_n [|F_{NH}(+) - F_{NH}(-)| - \Delta H_{cal}]^2 / n. \quad (2.4.4.31)$$

If all errors are assumed to reside in F_H , E can be evaluated in yet another way using the expression

$$E^2 = \sum_n (F_{HLE} - F_H)^2 / n. \quad (2.4.4.32)$$

2.4.4.7. Suggested modifications to Blow and Crick formulation and the inclusion of phase information from other sources

Modifications to the Blow and Crick procedure of phase evaluation have been suggested by several workers, although none represent a fundamental departure from the essential features of their formulation. In one of the modifications (Cullis, Muirhead, Perutz, Rossmann & North, 1961a; Ashida, 1976), all E_i 's are assumed to be the same, but the lack-of-closure error ξ_{Hi} for the i th derivative is measured as the distance from the mean of all intersections between phase circles to the point of intersection of the phase circle of that derivative with the phase circle of the native protein. Alternatively, individual values of E_i are retained, but the lack of closure is measured from the weighted mean of all intersections (Ashida, 1976). This is obviously designed to undo the effects of the unduly high weight given to F_N in the Blow and Crick formulation. In another modification (Raiz & Andreeva, 1970; Einstein, 1977), suggested for the same purpose, the F_N and F_{NH} circles are treated as circular bands, the width of each band being related to the error in the appropriate structure factor. A comprehensive set of modifications suggested by Green (1979) treats different types of errors separately. In particular, errors arising from imperfect isomorphism are treated in a comprehensive manner.

Although the isomorphous replacement method still remains the method of choice for the *ab initio* determination of protein structures, additional items of phase information from other sources are increasingly being used to replace, supplement, or extend the information obtained through the application of the isomorphous replacement. Methods have been developed for the routine refinement of protein structures (Watenpaugh, Sieker, Herriot & Jensen, 1973; Huber *et al.*, 1974; Sussmann, Holbrook, Church & Kim, 1977; Jack & Levitt, 1978; Isaacs & Agarwal, 1978; Hendrickson & Konert, 1980) and they provide a rich source of phase information. However, the nature of the problem and the inherent limitations of the Fourier technique are such that the possibility of refinement yielding misleading results exists (Vijayan, 1980a, b). It is therefore sometimes desirable to combine the phases obtained during refinement with the original isomorphous replacement phases. The other sources of phase information include molecular replacement (see Chapter 2.3), direct methods (Hendrickson & Karle, 1973; Sayre, 1974; de Rango, Mauguen & Tsoucaris, 1975) and different types of electron-density modifications (Hoppe & Gassmann, 1968; Collins, 1975; Schevitz, Podjarny, Zwick, Hughes & Sigler, 1981; Bhat & Blow, 1982; Agard & Stroud, 1982; Cannillo, Oberti & Ungaretti, 1983; Raghavan & Tulinsky, 1979; Wang, 1985).

The problem of combining isomorphous replacement phases with those obtained by other methods was first addressed by Rossmann & Blow (1961). The problem was subsequently examined by Hendrickson & Lattman (1970) and their method, which involves a modification of the Blow and Crick formulation, is perhaps the most widely used for combining phase information from different sources.

The Blow and Crick procedure is based on an assumed Gaussian 'lumped' error in F_{NH} which leads to a lack of closure, $\xi_{Hi}(\alpha)$, in F_{NH} defined by (2.4.4.20). Hendrickson and Lattman make an equally legitimate assumption that the lumped error, again assumed to be Gaussian, is associated with F_{NH} . Then, as in (2.4.4.20), we have

$$\xi_{Hi}''(\alpha) = F_{NH}^2 - D_{Hi}^2(\alpha), \quad (2.4.4.33)$$

where $\xi_{Hi}''(\alpha)$ is the lack of closure associated with F_{NH}^2 for an assumed protein phase angle α . Then the probability for

2.4. ISOMORPHOUS REPLACEMENT AND ANOMALOUS SCATTERING

α being the correct phase angle can be expressed as

$$P_i(\alpha) = N_i \exp[-\xi_{Hi}^2(\alpha)/2E_i^2], \quad (2.4.4.34)$$

where E_i^2 is the r.m.s. error in $F_{NH_i}^2$, which can be evaluated using methods similar to those employed for evaluating E_i . Hendrickson and Lautman have shown that the exponent in the probability expression (2.4.4.34) can be readily expressed as a linear combination of five terms in the following manner.

$$-\xi_{Hi}^2(\alpha)/2E_i^2 = K_i + A_i \cos \alpha + B_i \sin \alpha + C_i \cos 2\alpha + D_i \sin 2\alpha, \quad (2.4.4.35)$$

where K_i, A_i, B_i, C_i , and D_i are constants dependent on F_N, F_{Hi}, F_{NH_i} , and E_i^2 . Thus, five constants are enough to store the complete probability distribution of any reflection. Expressions for the five constants have been derived for phase information from anomalous scattering, tangent formula, partial structure, and molecular replacement. The combination of the phase information from all sources can then be achieved by simply taking the total value of each constant. Thus, the total probability of the protein phase angle being α is given by

$$P(\alpha) = \prod_i P_i(\alpha) = N \exp\left(\sum_s K_s + \sum_s A_s \cos \alpha + \sum_s B_s \sin \alpha + \sum_s C_s \cos 2\alpha + \sum_s D_s \sin 2\alpha\right), \quad (2.4.4.36)$$

where K_s, A_s , etc., are the constants appropriate for the sth source and N is the normalization constant.

2.4.4.8. Fourier representation of anomalous scatterers

It is often useful to have a Fourier representation of only the anomalous scatterers in a protein. The imaginary component of the electron-density distribution obviously provides such a representation. When the structure is known and $F_N(+)$ and $F_N(-)$ have been experimentally determined, Chacko & Srinivasan (1970) have shown that this representation is obtained in a Fourier synthesis with $i[F_N(+)+F_N^*(-)]/2$ as coefficients, where $F_N^*(-)$, whose magnitude is $F_N(-)$, is the complex conjugate of $F_N(+)$. They also indicated a method for calculating the phase angles of $F_N(+)$ and $F_N^*(-)$. It has been shown (Hendrickson & Sheriff, 1987) that the Bijvoet-difference Fourier synthesis proposed earlier by Kraut (1968) is an approximation of the true imaginary component of the electron density. The imaginary synthesis can be useful in identifying minor anomalous-scattering centres when the major centres are known and also in providing an independent check on the locations of anomalous scatterers and in distinguishing between anomalous scatterers with nearly equal atomic numbers (Sheriff & Hendrickson, 1987; Kitagawa, Tanaka, Hata, Katsube & Satow, 1987).

2.4.5. Anomalous scattering of neutrons and synchrotron radiation. The multiwavelength method

The multiwavelength anomalous-scattering method (Ramaseshan, 1982) relies on the variation of dispersion correction terms as a function of the wavelength used. The success of the method therefore depends upon the size of the correction terms and the availability of incident beams of comparable intensities at different appropriate wavelengths. Thus, although this method was used as early as 1957 (Ramaseshan, Venkatesan & Mani, 1957) as an aid to structure solution employing characteristic X-rays, it is, as outlined below,

ideally suited in structural work employing neutrons and synchrotron radiation. In principle, γ -radiation can also be used for phase determination (Raghavan, 1961; Moon, 1961) as the anomalous-scattering effects in γ -ray scattering could be very large; the wavelength is also easily tunable. However, the intensity obtainable for γ -rays is several orders lower than that obtainable from X-ray and neutron sources, and hence γ -ray anomalous scattering is of hardly any practical value in structural analysis.

2.4.5.1. Neutron anomalous scattering

Apart from the limitations introduced by experimental factors, such as the need for large crystals and the comparatively low flux of neutron beams, there are two fundamental reasons why neutrons are less suitable than X-rays for the *ab initio* determination of crystal structures. First, the neutron scattering lengths of different nuclei have comparable magnitudes whereas the atomic form factors for X-rays vary by two orders of magnitude. Therefore, Patterson techniques and the related heavy-atom method are much less suitable for use with neutron diffraction data than with X-ray data. Secondly, neutron scattering lengths could be positive or negative and hence, in general, the positivity criterion (Karle & Hauptmann, 1950) or the squarability criterion (Sayre, 1952) does not hold good for nuclear density. Therefore, the direct methods based on these criteria are not strictly applicable to structure analysis using neutron data, although it has been demonstrated that these methods could be successfully used in favourable situations in neutron crystallography (Sikka, 1969). The anomalous-scattering method is, however, in principle more powerful in the neutron case than in the X-ray case for *ab initio* structure determination.

Thermal neutrons are scattered anomalously at appropriate wavelengths by several nuclei. In a manner analogous to (2.4.3.1), the neutron scattering length of these nuclei can be written as

$$b_0 + b' + ib'' = b + ib''. \quad (2.4.5.1)$$

The correction terms b' and b'' are strongly wavelength dependent. In favourable cases, b'/b_0 and b''/b_0 can be of the order of 10 whereas they are small fractions in X-ray anomalous scattering. In view of this pronounced anomalous effect in neutron scattering, Ramaseshan (1966) suggested that it could be used for structure solution. Subsequently, Singh & Ramaseshan (1968) proposed a two-wavelength method for unique structure analysis using neutron diffraction. The first part of the method is the determination of the positions of the anomalous scatterers from the estimated values of F_Q . The method employed for estimating F_Q is analogous to that using (2.4.4.9) except that data collected at two appropriate wavelengths are used instead of those from two isomorphous crystals. The second stage of the two-wavelength method involves phase evaluation. Referring to Fig. 2.4.3.2 and in a manner analogous to (2.4.3.5), we have

$$\sin \psi_1 = \frac{F_{N1}^2(+)-F_{N1}^2(-)}{4F_{N1}F_{Q1}'}, \quad (2.4.5.2)$$

where $\psi = \alpha_N - \alpha_Q$ and subscript 1 refers to data collected at wavelength λ_1 . Singh and Ramaseshan showed that $\cos \psi_1$ can also be determined when data are available at wavelength λ_1 and λ_2 . We may define

$$F_m^2 = [F_N^2(+)+F_N^2(-)]/2 \quad (2.4.5.3)$$

and we have from (2.4.3.3), (2.4.3.4), and (2.4.5.3)

$$F_N = \{F_m^2 - F_Q^2\}^{1/2}. \quad (2.4.5.4)$$

2. RECIPROCAL SPACE IN CRYSTAL-STRUCTURE DETERMINATION

Then,

$$\cos \psi_1 = \frac{F_{m_1}^2 - F_{m_2}^2 - \{(b_1^2 + b_1'^2) - (b_2^2 + b_2'^2)\}x^2}{2(b_1 - b_2)F_{N_1}x} + \frac{F_{Q_1}}{F_{N_1}}, \quad (2.4.5.5)$$

where x is the magnitude of the temperature-corrected geometrical part of F_Q , ψ_1 and hence α_{N_1} can be calculated using (2.4.5.2) and (2.4.5.5). α_{N_2} can also be obtained in a similar manner.

During the decade that followed Ramaseshan's suggestion, neutron anomalous scattering was used to solve half a dozen crystal structures, employing the multiple-wavelength methods as well as the methods developed for structure determination using X-ray anomalous scattering (Koetzle & Hamilton, 1975; Sikka & Rajagopal, 1975; Flook, Freeman & Scudder, 1977). It has also been demonstrated that measurable Bijvoet differences could be obtained, in favourable situations, in neutron diffraction patterns from protein crystals (Schoenborn, 1975). However, despite the early promise held out by neutron anomalous scattering, the method has not been as successful as might have been hoped. In addition to the need for large crystals, the main problem with using this method appears to be the time and expense involved in data collection (Koetzle & Hamilton, 1975).

2.4.5.2. Anomalous scattering of synchrotron radiation

The most significant development in recent years in relation to anomalous scattering of X-rays has been the advent of synchrotron radiation (Helliwell, 1984). The advantage of using synchrotron radiation for making anomalous-scattering measurements essentially arises out of the tunability of the wavelength. Unlike the characteristic radiation from conven-

tional X-ray sources, synchrotron radiation has a smooth spectrum and the wavelength to be used can be finely selected. Accurate measurements have shown that values in the neighbourhood of 30 electrons could be obtained in favourable cases for f' and f'' (Templeton, Templeton, Phillips & Hodgson, 1980; Templeton, Templeton & Phizackerley, 1980; Templeton, Templeton, Phizackerley & Hodgson, 1982). Schemes for the optimization of the wavelengths to be used have also been suggested (Narayan & Ramaseshan, 1981). Interestingly, the anomalous differences obtainable using synchrotron radiation are comparable in magnitude to the isomorphous differences normally encountered in protein crystallography. Thus, the use of anomalous scattering at several wavelengths would obviously eliminate the need for employing many heavy-atom derivatives. The application of anomalous scattering of synchrotron radiation for macromolecular structure analysis is still in its infancy. However, the results obtained so far in this area have been encouraging (Helliwell, 1985). Intensity measurements from macromolecular X-ray diffraction patterns using synchrotron radiation have so far primarily relied upon oscillation photography (Arndt & Wonacott, 1977). This method is not particularly suitable for accurately evaluating anomalous differences. Much higher levels of accuracy are expected to be achieved with the use of area detectors (Arndt, 1986). Anomalous scattering, in combination with area detectors, is expected to develop into a major tool in macromolecular crystallography.

Acknowledgement

One of us (MV) acknowledges the support of the Department of Science & Technology, India.

References

- ADAMS, M. J. (1968). D. Phil. Thesis. Oxford Univ., England.
- AGARD, D. A. & STROUD, R. M. (1982). α -Bungarotoxin structure revealed by a rapid method for averaging electron density of non-crystallographically translationally related molecules. *Acta Cryst.* **A38**, 186-194.
- ARNDT, U. W. (1986). X-ray position-sensitive detectors. *J. Appl. Cryst.* **19**, 145-163.
- ARNDT, U. W. & WONACOTT, A. J. (1977). *The rotation method in crystallography*, Amsterdam: North-Holland.
- ASHIDA, T. (1976). Some remarks on the phase angle determination by the isomorphous replacement method. In *Crystallographic computing techniques*, edited by F. R. AHMED, pp. 282-284. Copenhagen: Munksgaard.
- BHAT, T. N. & BLOW, D. M. (1982). A density-modification method for improvement of poorly resolved protein electron-density maps. *Acta Cryst.* **A38**, 21-29.
- BIJVOET, J. M. (1949). Phase determination in direct Fourier-synthesis of crystal structures. *Proc. K. Ned. Akad. Wet. (B)*, **52**, 313-314.
- BIJVOET, J. M. (1954). Structure of optically active compounds in the solid state. *Nature (London)*, **173**, 888-891.
- BIJVOET, J. M., PEERDEMAN, A. F. & VAN BOMMEL, A. J. (1951). Determination of the absolute configuration of optically active compounds by means of X-rays. *Nature (London)*, **168**, 271-272.
- BLOW, D. M. & CRICK, F. H. C. (1959). The treatment of errors in the isomorphous replacement method. *Acta Cryst.* **12**, 794-802.
- BLOW, D. M. & MATTHEWS, B. W. (1973). Parameter refinement in the multiple isomorphous-replacement method. *Acta Cryst.* **A29**, 56-62.
- BLOW, D. M. & ROSSMANN, M. G. (1961). The single isomorphous replacement method. *Acta Cryst.* **14**, 1195-1202.
- BLUNDELL, T. L. & JOHNSON, L. N. (1976). *Protein crystallography*. London: Academic Press.
- BOKHOVEN, C., SCHOONE, J. C. & BIJVOET, J. M. (1951). The Fourier synthesis of the crystal structure of strychnine sulphate pentahydrate. *Acta Cryst.* **4**, 275-280.
- BRADLEY, A. J. & RODGERS, J. W. (1934). The crystal structure of the Heusler alloys. *Proc. R. Soc. London Ser. A*, **144**, 340-359.
- CANNILLO, E., OBERTI, R. & UNGARETTI, L. (1983). Phase extension and refinement by density modification in protein crystallography. *Acta Cryst.* **A39**, 68-74.
- CHACKO, K. K. & SRINIVASAN, R. (1970). On the Fourier refinement of anomalous dispersion corrections in X-ray diffraction data. *Z. Kristallogr.* **131**, 88-94.
- COLLINS, D. M. (1975). Efficiency in Fourier phase refinement for protein crystal structures. *Acta Cryst.* **A31**, 388-389.
- CORK, J. M. (1927). The crystal structure of some of the alums. *Philos. Mag.* **4**, 688-698.
- COSTER, D., KNOL, K. S. & PRINS, J. A. (1930). Unterschiede in der Intensität der Röntgenstrahlenreflexion an den beiden 111-Flächen der Zinkblende. *Z. Phys.* **63**, 345-369.
- CROMER, D. T. (1965). Anomalous dispersion corrections computed from self-consistent field relativistic Dirac-Slater wave functions. *Acta Cryst.* **18**, 17-23.

REFERENCES

- CRUICKSHANK, D. W. J. & McDONALD, W. S. (1967). Parameter errors in polar space groups caused by neglect of anomalous scattering. *Acta Cryst.* **23**, 9–11.
- CULLIS, A. F., MUIRHEAD, H., PERUTZ, M. F., ROSSMANN, M. G. & NORTH, A. C. T. (1961a). The structure of haemoglobin. VIII. A three-dimensional Fourier synthesis at 5.5 Å resolution: determination of the phase angles. *Proc. R. Soc. London Ser. A*, **265**, 15–38.
- CULLIS, A. F., MUIRHEAD, H., PERUTZ, M. F., ROSSMANN, M. G. & NORTH, A. C. T. (1961b). The structure of haemoglobin. IX. A three-dimensional Fourier synthesis at 5.5 Å resolution: description of the structure. *Proc. R. Soc. London Ser. A*, **265**, 161–187.
- DALE, D., HODGKIN, D. C. & VENKATESAN, K. (1963). The determination of the crystal structure of Factor V 1a. In *Crystallography and crystal perfection*, edited by G. N. RAMACHANDRAN, pp. 237–242. New York and London: Academic Press.
- DICKERSON, R. E., KENDREW, J. C. & STRANDBERG, B. E. (1961). The crystal structure of myoglobin: phase determination to a resolution of 2 Å by the method of isomorphous replacement. *Acta Cryst.* **14**, 1188–1195.
- DICKERSON, R. E., WEINZIERL, J. E. & PALMER, R. A. (1968). A least-squares refinement method for isomorphous replacement. *Acta Cryst.* **B24**, 997–1003.
- DODSON, E., EVANS, P. & FRENCH, S. (1975). The use of anomalous scattering in refining heavy atom parameters in proteins. In *Anomalous scattering*, edited by S. RAMASESHAN & S. C. ABRAHAMS, pp. 423–436. Copenhagen: Munksgaard.
- DODSON, E. & VJAYAN, M. (1971). The determination and refinement of heavy-atom parameters in protein heavy-atom derivatives. Some model calculations using acentric reflexions. *Acta Cryst.* **B27**, 2402–2411.
- EINSTEIN, J. E. (1977). An improved method for combining isomorphous replacement and anomalous scattering diffraction data for macromolecular crystals. *Acta Cryst.* **A33**, 75–85.
- FLOOK, R. J., FREEMAN, H. C. & SCUDDER, M. L. (1977). An X-ray and neutron diffraction study of aqua(L-glutamato)cadmium(II) hydrate. *Acta Cryst.* **B33**, 801–809.
- GILLI, G. & CRUICKSHANK, D. W. J. (1973). Effect of neglect of dispersion in centrosymmetric structures: results for OsO₄. *Acta Cryst.* **B29**, 1983–1985.
- GREEN, D. W., INGRAM, V. M. & PERUTZ, M. F. (1954). The structure of haemoglobin. IV. Sign determination by the isomorphous replacement method. *Proc. R. Soc. London Ser. A*, **225**, 287–307.
- GREEN, E. A. (1979). A new statistical model for describing errors in isomorphous replacement data: the case of one derivative. *Acta Cryst.* **A35**, 351–359.
- HARKER, D. (1956). The determination of the phases of the structure factors of non-centrosymmetric crystals by the method of double isomorphous replacement. *Acta Cryst.* **9**, 1–9.
- HELLIWELL, J. R. (1984). Synchrotron X-radiation protein crystallography: instrumentation, methods and applications. *Rep. Prog. Phys.* **47**, 1403–1497.
- HELLIWELL, J. R. (1985). Protein crystallography with synchrotron radiation. *J. Mol. Struct.* **130**, 63–91.
- HENDRICKSON, W. A. (1979). Phase information from anomalous-scattering measurements. *Acta Cryst.* **A35**, 245–247.
- HENDRICKSON, W. A. & KARLE, J. (1973). Carp muscle calcium-binding protein. III. Phase refinement using the tangent formula. *J. Biol. Chem.* **248**, 3327–3334.
- HENDRICKSON, W. A. & KONNERT, J. H. (1980). Incorporation of stereochemical information into crystallographic refinement. In *Computing in crystallography*, edited by R. DIAMOND, S. RAMASESHAN & K. VENKATESAN, pp. 13.01–13.23. Bangalore: Indian Academy of Sciences.
- HENDRICKSON, W. A. & LAITMAN, E. E. (1970). Representation of phase probability distributions in simplified combinations of independent phase information. *Acta Cryst.* **B26**, 136–143.
- HENDRICKSON, W. A. & SHERIFF, S. (1987). General density function corresponding to X-ray diffraction with anomalous scattering included. *Acta Cryst.* **A43**, 121–125.
- HENDRICKSON, W. A. & TEETER, M. M. (1981). Structure of the hydrophobic protein crambin determined directly from the anomalous scattering of sulphur. *Nature (London)*, **290**, 107–113.
- HOPPE, W. & GASSMANN, J. (1968). Phase correction, a new method to solve partially known structures. *Acta Cryst.* **B24**, 97–107.
- HUBER, R., KUKLA, D., BODE, W., SCHWAGER, P., BARTELS, K., DEISENHOFER, J. & STEIGENMANN, W. (1974). Structure of the complex formed by bovine trypsin and bovine pancreatic trypsin inhibitor. II. Crystallographic refinement at 1.9 Å resolution. *J. Mol. Biol.* **89**, 73–101.
- IBERS, J. A. & HAMILTON, W. C. (1964). Dispersion corrections and crystal structure refinements. *Acta Cryst.* **17**, 781–782. *International Tables for X-ray crystallography* (1974). Vol. IV, pp. 148–151. Birmingham: Kynoch Press. (Present distributor Kluwer Academic Publishers, Dordrecht.)
- ISAACS, N. W. & AGARWAL, R. C. (1978). Experience with fast Fourier least squares in the refinement of the crystal structure of rhombohedral 2-zinc insulin at 1.5 Å resolution. *Acta Cryst.* **A34**, 782–791.
- JACK, A. & LEVITT, M. (1978). Refinement of large structures by simultaneous minimization of energy and R factor. *Acta Cryst.* **A34**, 931–935.
- KARLE, J. & HAUPTMAN, H. (1950). The phases and magnitudes of the structure factors. *Acta Cryst.* **3**, 181–187.
- KARTHA, G. (1961). Isomorphous replacement in non-centrosymmetric structures. *Acta Cryst.* **14**, 680–686.
- KARTHA, G. (1965). Combination of multiple isomorphous replacement and anomalous dispersion data for protein structure determination. III. Refinement of heavy atom positions by the least-squares method. *Acta Cryst.* **19**, 883–885.
- KARTHA, G. (1975). Application of anomalous scattering studies in protein structure analysis. In *Anomalous scattering*, edited by S. RAMASESHAN & S. C. ABRAHAMS, pp. 363–392. Copenhagen: Munksgaard.
- KARTHA, G. (1976). Protein phase evaluation: multiple isomorphous series and anomalous scattering methods. In *Crystallographic computing techniques*, edited by F. R. AHMED, pp. 269–281. Copenhagen: Munksgaard.
- KARTHA, G. & PARTHASARATHY, R. (1965). Combination of multiple isomorphous replacement and anomalous dispersion data for protein structure determination. I. Determination of heavy-atom positions in protein derivatives. *Acta Cryst.* **18**, 745–749.
- KENDREW, J. C., DICKERSON, R. E., STRANDBERG, B. E., HART, R. G., PHILLIPS, D. C. & SHORE, V. C. (1960). Structure of myoglobin. A three-dimensional Fourier synthesis at 2 Å resolution. *Nature (London)*, **185**, 422–427.
- KITAGAWA, Y., TANAKA, N., HATA, Y., KATSUBE, Y. & SATOW, Y. (1987). Distinction between Cu²⁺ and Zn²⁺ ions in a crystal of spinach superoxide dismutase by use of anomalous dispersion and tuneable synchrotron radiation. *Acta Cryst.* **B43**, 272–275.

2. RECIPROCAL SPACE IN CRYSTAL-STRUCTURE DETERMINATION

- KOETZLE, T. F. & HAMILTON, W. C. (1975). Neutron diffraction study of NaSmEDTA.8H₂O: an evaluation of methods of phase determination based on three-wavelength anomalous dispersion data. In *Anomalous scattering*, edited by S. RAMASESHAN & S. C. ABRAHAMS, pp. 489-502, Copenhagen: Munksgaard.
- KRAUT, J. (1968). Bijvoet-difference Fourier function. *J. Mol. Biol.* **35**, 511-512.
- KRAUT, J., SIEKER, L. C., HIGH, D. F. & FREER, S. T. (1962). Chymotrypsin: a three-dimensional Fourier synthesis at 5 Å resolution. *Proc. Natl Acad. Sci. USA*, **48**, 1417-1424.
- MCPHERSON, A. (1982). *Preparation and analysis of protein crystals*. New York: John Wiley.
- MARK, H. & SZILLARD, L. (1925). Ein einfacher versuch zur auffindung eines selectiven effektes bei der zerstreung von Röntgenstrahlen. *Z. Phys.* **33**, 688-691.
- MATTHEWS, B. W. (1966). The determination of the position of anomalously scattering heavy atom groups in protein crystals. *Acta Cryst.* **20**, 230-239.
- MOON, P. B. (1961). Developments in gamma-ray optics. *Nature (London)*, **185**, 427-429.
- MUIRHEAD, H., COX, J. M., MAZZARELLA, L. & PERUTZ, M. F. (1967). Structure and function of haemoglobin. III. A three-dimensional Fourier synthesis of human deoxyhaemoglobin at 5.5 Å resolution. *J. Mol. Biol.* **28**, 156-177.
- NARAYAN, R. & RAMASESHAN, S. (1981). Optimum choice of wavelengths in the anomalous scattering technique with synchrotron radiation. *Acta Cryst.* **A37**, 636-641.
- NAVIA, M. A. & SIGLER, P. B. (1974). The application of direct methods to the analysis of heavy-atom derivatives. *Acta Cryst.* **A30**, 706-712.
- NORTH, A. C. T. (1965). The combination of isomorphous replacement and anomalous scattering data in phase determination of non-centrosymmetric reflexions. *Acta Cryst.* **18**, 212-216.
- OKAYA, Y. & PEPINSKY, R. (1956). New formulations and solution of the phase problem in X-ray analysis for noncentric crystals containing anomalous scatterers. *Phys. Rev.* **103**, 1645-1657.
- PARTHASARATHY, S. (1967). Expectation value of the Bijvoet ratio. *Acta Cryst.* **22**, 98-103.
- PARTHASARATHY, S. & SRINIVASAN, R. (1964). The probability distribution of Bijvoet differences. *Acta Cryst.* **17**, 1400-1407.
- PATTERSON, A. L. (1963). Treatment of anomalous dispersion in X-ray diffraction data. *Acta Cryst.* **16**, 1255-1256.
- PEERDEMAN, A. F. & BIJVOET, J. M. (1956). The indexing of reflexions in investigations involving the use of the anomalous scattering effect. *Acta Cryst.* **9**, 1012-1015.
- PEERDEMAN, A. F., VAN BOMMEL, A. J. & BIJVOET, J. M. (1951). Determination of the absolute configuration of optically active compounds by means of X-rays. *Proc. K. Ned. Akad. Wet. (B)*, **54**, 16-19.
- PEPINSKY, R., OKAYA, Y. & TAKEUCHI, Y. (1957). Theory and application of the $P_s(u)$ function and anomalous dispersion in direct determination of structures and absolute configuration in non-centric crystals. *Acta Cryst.* **10**, 756.
- PHILLIPS, D. C. (1966). Advances in protein crystallography. In *Advances in structure research by diffraction methods*, Vol. 2, edited by R. BRILL & R. MASON, pp. 75-140. New York and London: Interscience.
- RAGHAVAN, N. V. & TULINSKY, A. (1979). The structure of α -chymotrypsin. II. Fourier phase refinement and extension of the dimeric structure at 1.8 Å resolution by density modification. *Acta Cryst.* **B35**, 1776-1785.
- RAGHAVAN, R. S. (1961). On the possibility of a new phase-determining method applying the Mössbauer effect. *Proc. Indian Acad. Sci. Sect. A*, **53**, 265-271.
- RAIZ, V. SH. & ANDREEVA, N. S. (1970). Determining the coefficients of the Fourier series of the electron density function of protein crystals. *Sov. Phys. Crystallogr.* **15**, 206-210.
- RAMACHANDRAN, G. N. & RAMAN, S. (1956). A new method for the structure analysis of non-centrosymmetric crystals. *Curr. Sci.* **25**, 348-351.
- RAMACHANDRAN, G. N. & RAMAN, S. (1959). Syntheses for the deconvolution of the Patterson function. Part I. General principles. *Acta Cryst.* **12**, 957-964.
- RAMACHANDRAN, G. N. & SRINIVASAN, R. (1970). *Fourier methods in crystallography*. New York: Wiley-Interscience.
- RAMASESHAN, S. (1963). The use of anomalous scattering in crystal structure analysis. In *Advanced methods of crystallography*, edited by G. N. RAMACHANDRAN, pp. 67-95. London and New York: Academic Press.
- RAMASESHAN, S. (1966). The use of anomalous scattering of neutrons in the solution of crystal structures containing large molecules. *Curr. Sci.* **35**, 87-91.
- RAMASESHAN, S. (1982). Use of the multi-wavelength anomalous scattering method in crystallography. In *Computational crystallography*, edited by D. SAYRE, pp. 201-208. Oxford: Clarendon Press.
- RAMASESHAN, S. & ABRAHAMS, S. C. (1975). *Anomalous scattering*. Copenhagen: Munksgaard.
- RAMASESHAN, S. & VENKATESAN, K. (1957). The use of anomalous scattering without phase change in crystal structure analysis. *Curr. Sci.* **26**, 352-353.
- RAMASESHAN, S., VENKATESAN, K. & MANI, N. V. (1957). The use of anomalous scattering for the determination of crystal structures - KMnO₄. *Proc. Indian Acad. Sci.* **46**, 95-111.
- RANGO, C. DE, MAUGUEN, Y. & TSOUCARIS, G. (1975). Use of high-order probability laws in phase refinement and extension of protein structures. *Acta Cryst.* **A31**, 227-233.
- ROBERTSON, J. M. (1936). An X-ray study of the phthalocyanines. Part II. Quantitative structure determination of the metal-free compound. *J. Chem. Soc.* pp. 1195-1209.
- ROBERTSON, J. M. & WOODWARD, I. (1937). An X-ray study of the phthalocyanines. Part III. Quantitative structure determination of nickel phthalocyanine. *J. Chem. Soc.* pp. 219-230.
- ROSSMANN, M. G. (1960). The accurate determination of the position and shape of heavy-atom replacement groups in proteins. *Acta Cryst.* **13**, 221-226.
- ROSSMANN, M. G. (1961). The position of anomalous scatterers in protein crystals. *Acta Cryst.* **14**, 383-388.
- ROSSMANN, M. G. & BLOW, D. M. (1961). The refinement of structures partially determined by the isomorphous replacement method. *Acta Cryst.* **14**, 641-647.
- SAYRE, D. (1952). The squaring method: a new method for phase determination. *Acta Cryst.* **5**, 60-65.
- SAYRE, D. (1974). Least-squares phase refinement. II. High-resolution phasing of a small protein. *Acta Cryst.* **A30**, 180-184.
- SCHÉVITZ, R. W., PODIARNY, A. D., ZWICK, M., HUGHES, J. J. & SIGLER, P. B. (1981). Improving and extending the phases of medium- and low-resolution macromolecular structure factors by density modification. *Acta Cryst.* **A37**, 669-677.
- SCHOENBORN, B. P. (1975). Phasing of neutron protein data by anomalous dispersion. In *Anomalous scattering*, edited by S. RAMASESHAN & S. C. ABRAHAMS, pp. 407-416, Copenhagen: Munksgaard.

REFERENCES

- SHERIFF, S. & HENDRICKSON, W. A. (1987). Location of iron and sulfur atoms in myohemerythrin from anomalous-scattering measurements. *Acta Cryst.* B43, 209-212.
- SIKKA, S. K. (1969). On the application of the symbolic addition procedure in the neutron diffraction structure determination. *Acta Cryst.* A25, 539-543.
- SIKKA, S. K. & RAJAGOPAL, H. (1975). Application of neutron anomalous dispersion in the structure determination of cadmium tartrate pentahydrate. In *Anomalous scattering*, edited by S. RAMASESHAN & S. C. ABRAHAMS, pp. 503-514. Copenhagen: Munksgaard.
- SINGH, A. K. & RAMASESHAN, S. (1966). The determination of heavy atom positions in protein derivatives. *Acta Cryst.* 21, 279-280.
- SINGH, A. K. & RAMASESHAN, S. (1968). The use of neutron anomalous scattering in crystal structure analysis. I. Non-centrosymmetric structures. *Acta Cryst.* B24, 35-39.
- SRINIVASAN, R. (1972). Applications of X-ray anomalous scattering in structural studies. In *Advances in structure research by diffraction methods*, Vol. 4, edited by W. HOPPE & R. MASON, pp. 105-197. Braunschweig: Freidr. Vieweg & Sohn; and Oxford: Pergamon Press.
- SRINIVASAN, R. & CHACKO, K. K. (1970). On the determination of phases of a noncentrosymmetric crystal by the anomalous dispersion method. *Z. Kristallogr.* 131, 29-39.
- SUSSMANN, J. L., HOLBROOK, S. R., CHURCH, G. M. & KIM, S.-H. (1977). A structure-factor least-squares refinement procedure for macromolecular structures using constrained and restrained parameters. *Acta Cryst.* A33, 800-804.
- TEMPLETON, D. H., TEMPLETON, L. K., PHILLIPS, J. C. & HODGSON, K. O. (1980). Anomalous scattering of X-rays by cesium and cobalt measured with synchrotron radiation. *Acta Cryst.* A36, 436-442.
- TEMPLETON, L. K., TEMPLETON, D. H. & PHIZACKERLEY, R. P. (1980). L_3 -edge anomalous scattering of X-rays by praseodymium and samarium. *J. Am. Chem. Soc.* 102, 1185-1186.
- TEMPLETON, L. K., TEMPLETON, D. H., PHIZACKERLEY, R. P. & HODGSON, K. O. (1982). L_3 -edge anomalous scattering by gadolinium and samarium measured at high resolution with synchrotron radiation. *Acta Cryst.* A38, 74-78.
- TERWILLIGER, T. C. & EISENBERG, D. (1983). Unbiased three-dimensional refinement of heavy-atom parameters by correlation of origin-removed Patterson functions. *Acta Cryst.* A39, 813-817.
- UEKI, T., ZALKIN, A. & TEMPLETON, D. H. (1966). Crystal structure of thorium nitrate pentahydrate by X-ray diffraction. *Acta Cryst.* 20, 836-841.
- VIJAYAN, M. (1980a). On the Fourier refinement of protein structures. *Acta Cryst.* A36, 295-298.
- VIJAYAN, M. (1980b). Phase evaluation and some aspects of the Fourier refinement of macromolecules. In *Computing in crystallography*, edited by R. DIAMOND, S. RAMASESHAN & K. VENKATESAN, pp. 19.01-19.25. Bangalore: Indian Academy of Sciences.
- VIJAYAN, M. (1981). X-ray analysis of 2Zn insulin: some crystallographic problems. In *Structural studies on molecules of biological interest*, edited by G. DODSON, J. P. GLUSKER & D. SAYRE, pp. 260-273. Oxford: Clarendon Press.
- VIJAYAN, M. (1987). Anomalous scattering methods. In *Direct methods, macromolecular crystallography and crystallographic statistics*, edited by H. SCHENK, A. J. C. WILSON & S. PARTHASARATHY, pp. 121-139. Singapore: World Scientific Publishing Co. Pte. Ltd.
- VOS, A. (1975). Anomalous scattering and chemical crystallography. In *Anomalous scattering*, edited by S. RAMASESHAN & S. C. ABRAHAMS, pp. 307-317. Copenhagen: Munksgaard.
- WANG, B. C. (1985). Resolution of phase ambiguity in macromolecular crystallography. *Methods Enzymol.* 115, 90-112.
- WATENPAUGH, K. D., SIEKER, L. C., HERRIOT, J. R. & JENSEN, L. H. (1973). Refinement of the model of a protein: rubredoxin at 1.5 Å resolution. *Acta Cryst.* B29, 943-956.
- WATENPAUGH, K. D., SIEKER, L. C. & JENSEN, L. H. (1975). Anomalous scattering in protein structure analysis. In *Anomalous scattering*, edited by S. RAMASESHAN & S. C. ABRAHAMS, pp. 393-405. Copenhagen: Munksgaard.
- WILSON, A. J. C. (1942). Determination of absolute from relative X-ray intensity data. *Nature (London)*, 150, 151-152.
- WILSON, A. J. C. (1975). Effect of neglect of dispersion on apparent scale and temperature parameters. In *Anomalous scattering*, edited by S. RAMASESHAN & S. C. ABRAHAMS, pp. 325-332. Copenhagen: Munksgaard.
- ZACHARIASEN, W. H. (1965). Dispersion in quartz. *Acta Cryst.* 18, 714-716.